

# Cervical Artificial Disc Replacement

## What Is an Artificial Disc Replacement?

Cervical artificial disc replacement (arthroplasty) is a surgical procedure that involves decompressing spinal cord and nerves in the cervical spine and inserting a device that preserves motion in the spine. While the artificial disc allows the spine to still move, it is not as good as your normal disc.

In an artificial disc replacement, the surgeon accesses the cervical spine through an incision in the neck. The spinal cord and nerves are decompressed by removing the entire disc and any bone pressing on the nerves. A device that preserves motion is inserted in place of the disc.

## Why Do I Need This Procedure?

An artificial disc replacement may be recommended as a surgical treatment option for patients with pain in the neck or arms. There may also be weakness or numbness which has not responded to conservative treatment measures (rest, physical therapy or medication).

Dr. Smith will take a number of factors into consideration before recommending an artificial disc, including the condition to be treated, your age, health and lifestyle and your anticipated level of activity following surgery. It is important to note that only a very specific problem can be treated with an artificial disc, and fusion procedures such as an ACDF are more common in most patients.

It is extremely important to understand that as we get older, we all develop “changes” in our spines but not all of us have pain or neurologic problems. Your MRI report may mention many of these changes, but not all of them may explain your pain or neurologic problems. Dr. Smith will discuss which of the changes are causing problems and which are not causing problems. As a result, surgery is not intended to make your spine look “perfect” but instead to only address the changes that are causing your problem.

## How Do I Prepare For This Surgery?

- 1) **Stop smoking.** If you smoke, try to stop before your surgery. People who smoke have more disc problems and neck/ arm pain than people who don't. The complications of surgery are lower and recovery is quicker in non-smokers.
- 2) **Stop certain medications.** If you are on blood thinners such as Coumadin (Warfarin), Plavix, Pradaxa, Xarelto, Eliquis, notify Dr. Smith and your primary



care physician. These medications will need to be stopped prior to surgery and you will need to remain off them for a period following surgery. You may continue taking Aspirin if you are on it before surgery.

- 3) **Weight loss.** If you are overweight, then weight loss before surgery may lower complications and improve your recovery. Discuss with Dr. Smith and your primary care physician if it is recommended for you to try weight loss before surgery and how to go about the weight loss.
- 4) **Ask for time off work.** You will need to be off work for at least 4 weeks following surgery. It may be longer in certain circumstances. Make arrangements with your employer. We understand the financial constraints of many patients and the need to return to work as soon as possible. But please understand that if you return too early this may impair your healing and limit the beneficial effects of surgery.
- 5) **Stop eating and drinking the night before surgery.** It is standard to stop all food and drink the midnight before your surgery, even if your surgery is not first thing in the morning. You may also be asked to stop certain medications as well. If you are allowed to take some of your medications, you may take them with small sips of water. Definitely no coffee or juices the morning of surgery.
- 6) **Be a little early to the hospital.** The hospital will call you the day before surgery to notify you when to arrive at the hospital. Please be a little early. Dr. Smith performs many surgeries in a day and so your surgery time may be moved earlier than scheduled. Please be near your phone the day before and the day of surgery in case you are called of scheduling changes. Unfortunately, your surgery may also be delayed and we will notify you of this as well.
- 7) **Ask someone to drive you home and be available after surgery.** This is often an outpatient procedure but sometimes patients are admitted overnight. Either way, when you are discharged you cannot drive yourself home. You need to make these arrangements ahead of time.

## How Is an Artificial Disc Replacement Performed?

### The Operation

The operation is performed with the patient positioned on his or her back. You are completely asleep for the procedure, which usually lasts for 1-2 hours.



The entire disc is removed from the front and the spinal cord and nerves are decompressed



A device is placed where the disc used to be that allows motion at the level

## **What Happens After Surgery??**

You will wake up from surgery in the operating room or recovery area. After about 45 minutes, your family will be allowed to see you. Often the pain you are experiencing before surgery will be better but that is not always the case and sometimes takes longer to improve. You may experience a new soreness and muscle spasms in the back of your neck. This will improve with time and controlled with pain medications and muscle relaxers. Weakness and numbness often take days to weeks to months to improve. This often is an outpatient procedure although occasionally patients are admitted overnight. You will be up and walking the day after surgery. You do not need a brace following this surgery because we want you to move your neck.

## **What Happens When I Get Home??**

You still will be sore following surgery but it is important to be active following, walking and leaving your house occasionally. It will be much harder to fully recover if you stay in bed or sit in chairs all day. However you must not be too active. You cannot drive a car while in pain or on narcotics. You may twist and turn your neck but don't overdo it. No lifting anything heavier than a gallon of milk. You must leave your bandage on the incision. You may shower the day following surgery but leave the bandage alone, do not remove the bandage and do not submerge the bandage under water. If your bandage comes off on its own, do not attempt to cover it with your own bandage at home. If you start to see any redness or drainage call our office. You must avoid constipation following surgery. Pain medications and muscle relaxers may make you constipated so start taking a stool softener. If after a few days you still have not had a bowel movement, you may need to take a laxative, enema or even a suppository. It sounds trivial, but prolonged constipation will make you feel awful quickly. You need to make an appointment to see Dr. Smith ~2 weeks after surgery. You will be asked to take Ibuprofen 600mg every 6 hours for 2 weeks following surgery. This is not to control the pain but instead to help prevent the artificial disc from unintentionally fusing.

If you have any questions after surgery, please call our office between 8:30 and 5pm. If you need medications, remember you need to come to our office to pick them up. We cannot legally call in narcotic or muscle relaxing medications to your pharmacy. So if you are getting low on these medications, call our office ahead of time between 8:30 and 5pm Monday through Friday so you can pick up a prescription. If you take your medication more than we prescribe, we will not refill it early.

## **How Long Will It Take Me To Recover?**

Dr. Smith will explain your recovery program. After your 2 week office visit Dr. Smith will discuss how you are progressing. At one month it is possible for you to return to work depending on what type of work and if it can be light duty desk work. You may be prescribed therapy after surgery.

## Are There Any Potential Risks Or Complications?

All treatment and outcome results are specific to the individual patient. Results may vary. Dr. Smith cannot guarantee pain or neurologic deficit improvement. It is important to understand the risks to surgery and we have listed some below. Additionally, there may be risks we have not listed.

### Risks:

- 1) **Blood loss.** Blood loss is usually very small. But as with any surgery, there is the potential for major or even life-threatening blood loss.
- 2) **Infection.** Even with antibiotics and careful technique, there is still a small risk of developing infection. This could require antibiotics or even further surgery to resolve. Infections may result in residual pain or neurologic deficits including weakness, sensory changes or bowel/bladder incontinence. Unfortunately this could become permanent.
- 3) **Reaction to anesthesia.** Anytime you are given medications you can experience an adverse reaction. Even if given medications you have tolerated in the past, you can develop new reactions.
- 4) **Cerebrospinal Fluid (CSF) leak.** Your nerves sit in a sac, which contains your nerves and spinal fluid. During surgery the sac may accidentally be punctured or opened. When Dr. Smith observes this he will attempt to fix it during surgery. However the fluid may still leak or Dr. Smith may not see it leaking. This may lead to headaches after surgery. At the surgery site a bump under your skin may occur or the fluid may even leak out of the incision. This could lead to infection or other problems requiring further surgery.
- 5) **Damage to the spinal cord or nerves.** The surgery is performed around your nerves. In the process of decompressing your spinal cord or nerves, injury can occur including pain, weakness, sensory changes or bowel/bladder incontinence. Unfortunately this could become permanent and severe. It may require additional surgery to improve.
- 6) **Fusion.** The purpose of surgery is to preserve your range of motion and avoid fusion, however rarely these devices still fuse.
- 7) **Loss of range of motion.** The device is intended to preserve your range of motion but it is not as good as your normal disc. You may still notice some loss of range of motion.
- 8) **Hematoma.** There is always bleeding during surgery and unfortunately a small amount of blood can collect and press into the nerves. This sometimes requires further surgery to decompress.
- 9) **Adjacent level disease.** Surgery attempts to preserve your range of motion and this theoretically avoids extra stress on the other spinal levels. However even with an artificial disc, some people develop arthritis at nearby spinal levels and sometimes require surgery on those levels.

- 10) **Hardware failure.** The device can fail, break or dislodge. There is a risk that the device could move and press into your nerves or spinal cord. There is a chance the device would need to be removed, replaced or converted to a fusion in the future.
- 11) **Dysphagia.** Almost everyone following surgery develop some swallowing trouble. Usually while eating hard foods such as breads, meat or pills the food “gets caught” around the area of the incision. This almost always improves with time but sometimes does not.
- 12) **Hoarseness.** Injury to the nerve that controls your voicebox could result in hoarseness. This could range from a little raspiness to a very soft voice. This almost always improves with time but sometimes does not.
- 13) **Failure to relieve symptoms.** Dr. Smith will do everything possible to give you the best results with the surgery. However, surgery may not relieve all or any of your symptoms.
- 14) **Reoperation.** As listed above, there are numerous scenarios which may require additional surgery in the future. Whether for reasons listed above or reasons not listed, undergoing surgery now does not preclude you from potentially needing surgery in the future.
- 15) **Death.** As with any surgical procedure, there is a risk of death. This is rare.